

intermak  
**VITALIC** **DiATOX**

**EFFECTIVE FEED ADDITIVE AGAINST EXTERNAL  
AND INTERNAL PARASITES**



DiaTOX is a natural product  
based on organic silicon. Lice,  
fleas, ticks, scabies, mites etc.

It ensures the elimination of  
external parasites. It is also a feed  
additive that supports digestion.






## Harms of Parasites

- Stress in the herd
- Decreased feather quality
- Skin irritation
- Weight loss
- Anemia
- Up to 20% decrease in egg production
- Decreased egg quality
- Increased death

Red mites can carry viruses and bacteria that cause disease in chickens (eg Salmonella, Pasteurella, Leukosis...)




## Mode of Action of DiaTOX on Parasites

Diatox, which consists of the fossilized siliceous shells of single-celled, microscopic algae, resembles sharp-edged broken glass pieces when viewed under a microscope, and with this feature, it destroys the waxy exoskeleton (cuticle) of the parasites, absorbs its moisture and kills by ~~dries~~ it from the inside.

by drying it

It acts physically, not chemically.



Scanning electron microscope (SEM) image

## Benefits

It ensures the elimination of external parasites like lice, fleas, ticks, scabies, mites.

- ~~Lice, fleas, ticks, scabies, mites and. It ensures the elimination of external parasites.~~
- It helps reduce bacterial infection.
- It reduces humidity and odor in poultry houses, corrals and shelters.
- It can also be used as a litter.
- Eliminates parasites in stored grains.

**DiaTOX is completely natural and does not contain any chemicals. Thanks to this feature, animals can safely consume;**

- It ensures the destruction of internal parasites such as worms.
- Supports the gut.
- It dries the feces and reduces fecal odor in poultry.
- Contains 15 trace minerals.
- Reduces feather loss caused by pecking.
- It aids digestion.

**THIS IS NOT A DRUG, IT IS A FEED ADDITIVE. IT IS NOT USED FOR TREATMENT PURPOSES.**



## Application

DiaTOX can be used for all poultry species.

**In the coops; 30-50 g/m<sup>2</sup>**

Apply the powder to the coop by spreading the powder under the wings, around the ventilation zone, in cracks, crevices and nest areas where mites hid

**In the poultry; 1 kg of feed per ton**

## Duration of Action

It shows its effect within 3 days. It may be necessary to repeat the application after 10 days. If the applied floor is covered with feces and feathers, the need for repetition ~~is formed.~~ **the need for repetition occurs.**

## Warnings and Precautions

- Mask should be worn during application.
- The product should be stored in a dry place that will not take moisture and should be applied to the dry surface.
- Do not use on animals other than the specified species.
- Please contact for information.